



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

DEPARTMENTS.

NOTE. All solutions of problems, problems for solution, and other department contributions should be sent direct to THE AMERICAN MATHEMATICAL MONTHLY, 1227 Clay Street, Springfield, Mo.

SOLUTIONS OF PROBLEMS.

NOTE. The following problems were received too late for publication: Calculus No. 200, and Geometry No. 263, solved by G. W. Greenwood. Credit is also given to J. Scheffer for solutions of Calculus No. 201, Diophantine Analysis No. 127, and Geometry No. 262.

ALGEBRA.

237. Proposed by F. P. MATZ, Sc. D., Ph. D., Reading, Pa.

Solve $x^2 + y + z = 12 \dots (1)$; $x + y^2 + z = 8 \dots (2)$; $x + y + z^2 = 6 \dots (3)$.

Solution by L. E. NEWCOMB, Los Gatos, Cal.

Since (3) from (1) gives $x^2 - x = 6 + z^2 - z$, $\therefore x = \frac{1}{2} \pm \sqrt{(6\frac{1}{4} + z^2 - z)} \dots (4)$;
(3) from (2) gives $y = \frac{1}{2} \pm \sqrt{(2\frac{1}{4} + z^2 - z)} \dots (5)$.

Substitute these values for x and y in (3); then $\frac{1}{2} + \sqrt{(6\frac{1}{4} + z^2 - z)} + \frac{1}{2} + \sqrt{(2\frac{1}{4} + z^2 - z)} + z^2 = 6$.

Whence $[\sqrt{(6\frac{1}{4} + z^2 - z)} + \sqrt{(2\frac{1}{4} + z^2 - z)}]^2 = (5 - z^2)^2 \dots (6)$.

After expansion and transposition, (6) becomes $2\sqrt{(6\frac{1}{4} + z^2 - z)}(2\frac{1}{4} + z^2 - z) = z^4 - 12z^3 + 2z + 16\frac{1}{2}$. Square both numbers; then $z^8 - 24z^6 + 4z^5 + 173z^4 - 40z^3 - 430z^2 + 100z + 216 = 0$. The roots are $z = +1$, $+ .204923$, $+ 2.39427$, $+ 3.48865$, $- 3.806118$, $- 2.36109$, $- 2.09446$, $- .67044$.

Similarly two equations, one involving x , the other, y , are derived; or the values of x , y may be found from (4), (5), respectively. These values are:

$x = 3$, $+ 3.3983$, $- 2.59649$, $- 3.3642$, $- 4.45405$, $- 3.2664$, $+ 4.06808$, $+ 3.21476$;
 $y = 2$, $- 1.5976$, $+ 2.86394$, $- 2.80636$, $- 4.0324$, $+ 3.69155$, $- 2.4548$, $+ 2.33574$.

Also solved by A. H. Holmes, J. Scheffer, G. B. M. Zerr, and the Proposer.

238. Proposed by S. A. COREY, Hiteman, Iowa.

Prove that $\frac{1}{n+1} + \frac{1}{3(n+3)} + \frac{1}{5(n+5)} + \dots = \frac{1}{2} \left[\frac{1}{(n-1)} + \frac{1}{3(n-3)} + \frac{1}{5(n-5)} + \dots + \frac{1}{l(n-l)} \right]$, n being an even positive integer and $l = n - 1$.

Solution by J. SCHEFFER, A. M., Hagerstown, Md.

Putting successively $m = 2, 4, 6, 8, \dots, -m$, we get

$\frac{1}{1.3} + \frac{1}{3.5} + \frac{1}{5.7} + \dots$ ad inf. $= \frac{1}{2} \left[\left(\frac{1}{1} - \frac{1}{3} \right) + \left(\frac{1}{3} - \frac{1}{5} \right) + \left(\frac{1}{5} - \frac{1}{7} \right) + \dots \right] = \frac{1}{2}(1) = \frac{1}{2}$.